

Summer-winter comparisons of oxygen, nutrients and carbonates in the polar seas*

Chen-Tung A. CHEN**

Abstract: Until very recently, marine chemistry studies in the polar seas were mainly performed in summer. Little or no chemical data were collected in winter because of operational problems. Yet, deep and bottom waters of the world oceans were mainly formed in winter in the polar regions. The lack of winter data in such regions has prevented scientists from accurately estimating the variation of chemical properties of deep and bottom waters as they move away from their formation regions.

Winter chemical data, including oxygen, nutrients, alkalinity, total CO₂, pH and P_{CO₂} have now been collected in the Weddell Sea, the South Indian Ocean, and the Bering Sea. The concentrations of certain chemicals, such as oxygen, were found to differ drastically from summer values. Some properties, such as alkalinity, on the other hand, do not show significant seasonal variations when compared at the same salinity and temperature.

1. Introduction

Subsurface waters are mainly formed in the high latitude oceans in the wintertime. One must therefore study the water chemistry in these regions in winter in order to understand the global biogeochemical cycle of chemicals. Owing to operational problems, however, few chemical measurements have been made in the high latitude regions in winter. For instance, we know of only seven oceanographic stations with high-precision carbonate data in the entire Indian Ocean south of 30°S. And all these stations were occupied in the austral summer (TAKAHASHI *et al.*, 1980). Not knowing the characteristic properties of the subsurface waters near their origin, and near the time of their formation, makes it difficult, if not impossible, to interpret variations in the carbonate chemistry in the Indian Ocean.

We recently had the opportunity to participate in winter cruises in the Weddell Sea, the South Indian Ocean, and in the Bering Sea. What

follows will be the comparison between our winter data and the summer data collected by us and by other investigators.

2. Study regions and measurements

a) SOMOV data: Concurrent pH and titration alkalinity (TA) were measured on the research/supply vessel, MIKHAIL SOMOV, of the Arctic-Antarctic Research Institute of Leningrad, USSR. The Weddell Polynya Expedition (WEPOLEX) started in Montevideo, Uruguay, on 9 October 1981, crossed the Polar Front, and entered the ice field at 5°E, 56°30'S. SOMOV reached its southernmost point within the ice field at 62°20'S, and left the ice edge at 57°30'S near the Greenwich Meridian after obtaining chemical data from 20 vertical stations. After crossing the Polar Front again, the ship returned to Montevideo on 25 November. The experimental technique and data are described and listed elsewhere (CHEN, 1982a; 1983; 1984). Supporting oxygen and nutrient data are described and listed in JENNINGS *et al.*, (1984), HUBER *et al.* (1983) and GORDON *et al.* (1984).

b) MARION DUFRESNE data: Concurrent pH, TA, total CO₂ (TCO₂) and nitrate (NO₃) data were obtained in the wintertime southwestern Indian Ocean as part of the INDIVAT 1 Expedition (INDIEN VALORISATION de

* Received June 2, 1987

** On leave from the College of Oceanography, Oregon State University, Corvallis, Oregon 97331, U.S.A.

Present address: Institute of Marine Geology, National Sun Yat-Sen University, Kaohsiung, Taiwan, Republic of China

TRANSIT) aboard the French research/supply vessel MARION DUFRESNE. The ship departed La Reunion on 3 July 1984, reoccupied the GEOSECS station GS427, crossed the Subtropical Front at about 40°S, stopped in Crozet, reoccupied GS429 after crossing the Polar Front, then proceeded to Kerguelen and Amsterdam after crossing the Polar and Subtropical Fronts again, and returned to La Reunion on 4 August. Concurrent pH, TA and TCO₂ data were again obtained in the same general region in the austral summer of 1985 as part of the INDIGO 1/INDIVAT 3 Expedition (INDIGO stands for INDIEN GAS OCEAN) aboard the MARION DUFRESNE. The vessel departed La Reunion on 23 February and returned on 30 March after collecting chemical data from 23 stations including four GEOSECS stations (GS427-429, 454). The Subtropical Front was crossed at approximately 43°S and the Antarctic Front near 52°S.

The data description and listing are given in CHEN and POISSON (1986) and CHEN *et al.* (1986). The supporting oxygen and nutrient data are given in POISSON *et al.* (in preparation).

c) POLAR SEA data: The POLAR SEA left Dutch Harbor, Alaska, on 18 February and returned on 19 March 1983, after moving in and out of the Bering Sea ice between 58° and 61°30'N and between 171° and 179°W. Concurrent pH, TA and oxygen concentrations were measured. The data have been described and listed in CHEN (1985) and CHEN *et al.* (1985).

Approximate study areas of these expeditions are shown in Fig. 1.

3. Comparison of the oxygen data

BROECKER *et al.* (1985) reported that the "surface ocean waters are almost always found to be supersaturated with dissolved oxygen," and that the amount of supersaturation averages about 7 μ mol/kg or approximately 3%. The available data contradict this conclusion. Although BROECKER *et al.* (1985) investigated three of the largest data sets available (those of GEOSECS, NORPAX, and TTO), they did not examine the reports, including those containing winter data, published for high-latitude regions. LEVITUS (1982) summarized all data from the National Oceanographic Data Center as of 1978 and published the annual and seasonal mean oxygen saturation levels at the sea surface. The all-data annual mean indeed shows surface supersaturation except in upwelling areas of the eastern equatorial Pacific and the Southern Ocean. LEVITUS (1982), however, cautioned against a hastened conclusion, because the all-data annual mean may represent data from only one season, most likely summer. The Southern Ocean annual mean, for example, was shown to be a few percent undersaturated, but at that time sufficient winter data were not available. The recent winter data, to be discussed later, indicate an even lower degree of saturation. The then available and recent winter data also




Fig. 1. Study regions of SOMOV, MARION DUFRESNE and POLAR SEA.
Map provided by A. Mantyla.




Fig. 2. Correlation between (a) temperature and oxygen, and (b) temperature and % oxygen saturation for Bering Sea shelf waters in winter (Taken from CHEN, 1985).

show undersaturation in the Bering Sea and the northern North Atlantic, but because the more abundant summer data show supersaturation, the annual all-data mean also shows supersaturation.

Clearly, not all surface ocean waters are supersaturated with oxygen, and certainly they are not always supersaturated. Analysis of oxygen saturation for the four seasonal periods is desirable, but we will only attempt to make the summer-winter comparison because of the limitation in data coverage.

As early as 1963, SMETANIN (1963) reported that wintertime northeast Pacific surface water is slightly undersaturated with oxygen.

IVANENKOV (1964) reported that nearly all of the Bering Sea in winter absorbs oxygen. He reported an average of 6% undersaturation in the western Bering Sea, a value later confirmed by REID (1973). A similar degree of oxygen undersaturation exists in the northwestern North Pacific Ocean in winter (HAKODATE MARINE OBSERVATORY, 1967; REID, 1973; 1982). The wintertime northern Gulf of Alaska surface water data and our POLAR SEA data on the eastern Bering Sea shelf also show systematic

undersaturation in winter (SIO Reference 70-5, 1970; CHEN, 1985; CHEN *et al.*, 1985). The oxygen content of the POLAR SEA winter surface layer is higher than that of the deep layer on the ice-covered shelf (Fig. 2a), but the degree of saturation (CHEN, 1981) is reversed, *i. e.* most of the deep shelf water is more highly saturated than that of the surface water (Fig. 2b). This phenomenon is not observed in summer when both the oxygen content and the degree of saturation of the surface water are higher than those of the deep shelf water (*e. g.* HATTORI, 1977; 1979).

In summer, shelf oxygen data show extreme variability, governed by biological processes (CODISPOTI *et al.*, 1986). Winter oxygen, however, seems to mix conservatively regionally as suggested by segments of linear temperature vs oxygen correlations (Fig. 2a), discussed in detail in CHEN (1985).

Waters cooler than -0.6°C are all in the homogeneous surface layer and have higher oxygen concentration at lower temperature (Fig. 2a). All waters below -0.6°C , however, remain at about 5% undersaturation (Fig. 2b), suggesting that air-sea exchange and photo-




Fig. 3. Correlation between (a) temperature and oxygen, (b) temperature and % oxygen saturation of our winter Aleutian Basin data, and correlation between (c) temperature and oxygen, (d) temperature and % oxygen saturation of the summer *Hakuho Maru* data from the same location. The broken line shows the theoretical slope (Taken from CHEN, 1985).

synthesis cannot replenish oxygen fast enough to compensate for upwelling of water with lower O₂ content, respiration, and cooling effects. The result is that the surface waters actually show a lower degree of oxygen saturation than most deep waters (CHEN, 1985; CHEN *et al.*, 1985).

Deep waters for POLAR SEA stations (104 and 105) in the Aleutian Basin show a minimum in oxygen (Fig. 3a). The homogeneous, ice-free surface layer is also undersaturated with respect to oxygen by about 8% (Fig. 3b). The winter surface water oxygen concentration is

similar to the value found in the minimum temperature layer in the summer. Summer data from the same locality (stations KH78-3-8 and KH78-3-9 of the *Hakuho Maru* Cruise; HATTORI, 1979) are plotted in Fig. 3c and 3d for comparison. Seemingly, summer warming raises the temperature, but the oxygen content of the surface layer does not increase by much except in the top thirty meters, where supersaturation is obviously caused by enhanced photosynthesis. In winter, enhanced vertical mixing and rapid cooling, unmatched by slower air-sea oxygen

exchange and photosynthesis, produce undersaturation in the surface layer (IVANENKOV, 1964; REID, 1973; CHEN, 1985). BRUJEWIEZ *et al.* (1960) also found an undersaturation as much as 20% in the summer minimum-temperature layer in the Sea of Okhotsk, with the lowest degree of saturation on the continental shelf off Siberia and Sakhalin Island where intensive cooling occurs in winter.

The situation is similar in the Weddell Sea, based on the comparison of the SOMOV (CHEN, 1982a; 1984; HUBER, *et al.*, 1983; GORDON, *et al.*, 1984), GEOSECS (TAKAHASHI *et al.*, 1980) and other data in the literature (BRENNECKE, 1921; DEACON, 1940; OB, 1958; 1959; SECRETARIA DE MARINA, 1959a, b, c; IGY, 1961; some of these data need to be adjusted due to systematic analytical error). An ice-free summertime GEOSECS Atlantic station, GS89, is located at 0.0°E, 60.0°S which is very close to the ice-covered SOMOV station 33 (0°20'E, 60°S). The θ/S plot for waters below the θ_{min} layer at GS89 is essentially the same as the plot at SOMOV 33. The θ/AOU plot (Fig. 4) below the θ_{min} layer at GS89 is also similar to the plot at SOMOV 33. The SOMOV 33 data show a high AOU (apparent oxygen utilization) value of 50 $\mu\text{mol/kg}$ at the surface because the surface water is mixed with low-oxygen Weddell Deep Water, while the ice blocked the




Fig. 4. θ/AOU plot for WEPOLEX station 33 and GEOSECS station 89 in the Weddell Sea. The temperature minimum layer for the GEOSECS data is at about -1.8°C .

input of atmospheric oxygen (GORDON *et al.*, 1984). This finding supports the suggestion of WEISS *et al.* (1979), EDMOND *et al.* (1979), MINAS (1980), LYAKHIN and RUSANOV (1980) and CHEN (1982b) but disagrees with the Arctic work of GOSINK *et al.* (1976) and F. HERR (private communication, 1983) who found that the annual Arctic sea ice is permeable to gases.

Our winter data in the Bering Sea also suggest some air-sea exchange, but the exchange could have happened in the numerous leads and polynyi that we encountered in the Bering Sea but not in the Weddell Sea. Furthermore, unlike the annual Arctic sea ice which has many brine channels, the Weddell Sea ice has a large content of frazil ice with few brine channels, resulting in low permeability (ACKLEY *et al.*, 1980; 1982; CLARKE and ACKLEY, 1982). The late-winter, early-spring Weddell Sea surface water is at about 86% saturation, higher than the 60% for deep waters (BRENNECKE, 1921; GORDON *et al.*, 1984). The winter SEAMUNDSSON and HUDSON data (MALMBERG, 1983; CSS HUDSON, 1984) in the northern North Atlantic Ocean also show systematic oxygen undersaturation (as low as 10% undersaturation) within or out of the ice field.

The above discussion clearly indicates that the high-latitude regions are probably undersaturated




Fig. 5. Surface normalized nitrate concentrations vs temperature for the GEOSECS, INDIVAT 1 and INDIGO 1/INDIVAT 3 data in the south Indian Ocean.

with oxygen in winter, although water chemistry beneath the ice varies from place to place, and we cannot yet generalize the findings in one area to represent other regions. As further note, ANDERSON and DYRSSEN (1980) reported surface oxygen supersaturation within the summer Barents Sea ice field, which is, no doubt, due to photosynthesis.

4. Comparison of the nutrient data

Surface nutrient values frequently show significant variations between cruises even when data from the same location are compared. For instance, at GS 429 occupied in summer (Feb. 1978), the surface nitrate concentration was $20.2 \mu\text{mol/kg}$ (WEISS *et al.*, 1983). The INDIGO 1/INDIVAT 3 value was $21.2 \mu\text{mol/kg}$ (March 1985) and the INDIVAT 1 value was $24.5 \mu\text{mol/kg}$ (July 1984). For unknown reasons, however, the surface nitrate value correlates linearly with temperature when the nitrate concentrations are normalized to a constant salinity basis. The normalized nitrate ($NNO_3 = NO_3 \times 35/S$) data from the above three cruises are shown in Fig. 5. These three temperature trends show much

smaller variations. Phosphate and silicate behave in a similar way. Parts of the natural variations, such as those due to evaporation and precipitation, are removed by normalization, thus the resulting trends reflect mainly biological and mixing processes.

5. Comparison of the alkalinity and total CO_2 data

The surface alkalinity and total CO_2 values show large seasonal variations when compared in the same region, even when the effect of precipitation and evaporation is considered. The potential temperature is plotted against normalized total CO_2 ($NT\text{CO}_2$) for the WEPOLEX and GEOSECS data in the Weddell Sea (Fig. 6). The WEPOLEX winter data agree with the GEOSECS summer data below the temperature minimum layer, which is the remnant winter surface water (CHEN, 1984; POISSON and CHEN, 1987). The summer surface water, however, is lower by as much as $50 \mu\text{mol/kg}$ in $NT\text{CO}_2$ because of biological consumption. The NTA values are approximately $10 \mu\text{eq/kg}$ lower.

The surface NTA and $NT\text{CO}_2$ values, how-




Fig. 6. Potential temperature vs normalized total CO_2 for the WEPOLEX (all data) and GEOSECS (stations 79, 82, 85, 87 and 89) expeditions in the Weddell Sea. The temperature minimum layer for the GEOSECS data is between 0 and -1.8°C where $NT\text{CO}_2$ approximately $2,250 \mu\text{mol kg}^{-1}$.




Fig. 7. Surface (a) normalized total CO_2 and (b) alkalinity vs temperature for the GEOSECS and SOMOV data in the south Atlantic Ocean.

ever, show less seasonal variation when compared at the same temperature. The normalized surface alkalinity and total CO_2 seem to decrease steadily with increasing temperature (CHEN and MILLERO, 1979; CHEN and PYTKOWICZ, 1979). The GEOSECS NTCO_2 data (Dec. 1972–Jan. 1973) in the southern South Atlantic Ocean are higher than the October SOMOV data by $6 \mu\text{mol/kg}$ and are significantly higher than the November SOMOV data above 5°C (Fig. 7a). These differences reflect a possible seasonal effect due to changes in biological productivity. The biological productivity, however, does not significantly affect alkalinity. Indeed we could not detect any seasonal variability in NTA vs temperature correlations among the GEOSECS, SOMOV October and SOMOV November data (Fig. 7b). The GEOSECS NTA data are approximately $5 \mu\text{eq/kg}$ higher than the SOMOV data, but the difference is slightly smaller than the combined experimental error.

Data in the south Indian Ocean also show strong seasonal variations in NTCO_2 when the GEOSECS data are compared with the INDI-




Fig. 8. Surface pH values (measured at 25°C) vs temperature of the SOMOV data (taken from CHEN, 1984) in the South Atlantic Ocean.

VAT 1 and INDIGO 1/INDIVAT 3 data. The difference in NTA is smaller, on the order of $10 \mu\text{eq/kg}$, only slightly larger than the combined analytical error.

6. Comparison of the pH and P_{CO_2} data

The surface pH values (measured at 25°C) also correlate linearly with temperature, and seasonal variations are evident from the SOMOV data (Fig. 8). MARION DUFRESNE data in the south Indian Ocean also show linear correlations with temperature but there the summer trend does not differ significantly from the winter trend below 18°C .

It is well known that pH variations in the subsurface waters follow variations in oxygen. Fig. 9 shows the pH vs temperature correlations of our POLAR SEA winter and *Hakuho Maru*'s 78-3 summer Aleutian Basin stations (HATTORI, 1979). The curves are of similar shape as the temperature vs oxygen plots (Figs. 3a, c). Our deep water values are systematically higher than the *Hakuho Maru* data by 0.05 pH units, probably owing partly to a difference in calibration. The winter surface value would agree with the summer value at the minimum temperature layer if the summer value is systematically shifted up 0.05 units. The summer surface pH value are




Fig. 9. Correlation between temperature and pH values (measured at 25°C) of (a) our winter Aleutian Basin data and (b) the summer *Hakuho Maru* data from the same location (Taken from CHEN, 1985).

significantly higher than the winter value, reflecting CO_2 consumption by primary productivity (CHEN, 1985).

PCO_2 values are also known to show large seasonal variations (DEACON, 1940). Strangely, our SOMOV data (CHEN, 1984) and the data of TAKAHASHI (private communication, 1982) and TAKAHASHI and CHIPMAN (1982) are similar to the GEOSECS data (TAKAHASHI *et al.*, 1980) near 60°S in the south Atlantic and are nearly in equilibrium with the atmosphere. Rapid cooling of the surface water in winter reduces PCO_2 , but upwelling and entrainment (GORDEN *et al.*, 1984) increase PCO_2 . It is fortuitous that the net result is near equilibrium with the atmosphere. These observations prompted the suggestion that the Antarctic surface waters are likely to be in equilibrium with the atmosphere throughout the year (CHEN, 1984; TAKAHASHI, private communication, 1982; TAKAHASHI and CHIPMAN, 1982). Recent observations of TAKAHASHI (private communication, 1985), however, do show large temporal variations in the Southern Ocean. Fig. 10, based on the SOMOV data,




Fig. 10. Surface PCO_2 values vs seawater temperature of the SOMOV data (taken from CHEN, 1984) in the south Atlantic Ocean.

also shows large October to November changes in PCO_2 at high temperatures.

7. Conclusion

Certain chemical properties such as oxygen, pH and PCO_2 show large seasonal variations in the polar seas. Nutrients, alkalinity, and total CO_2 also vary significantly at the same location. But the variation is much reduced when compared at the same salinity and temperature. The seasonal variability in the normalized alkalinity is not much higher than the analytical error.

We believe that winter values should be used in preference to the summer values when one needs to know the initial, or preformed, concentrations of chemicals for deep and bottom waters. Without such information, calculations of production or consumption rates of chemicals are frequently subject to large error.

Acknowledgment

Funding for participation in the SOMOV, POLAR SEA and MARION DUFRESNE cruises and for data analysis was provided by the U.S. Department of Energy (81 EV 10611, A001, A002 and 19X-89608C under Martin Marietta Energy Systems, Inc., contract DE-AC 05-84 OR21440), the U.S. National Science Foundation (OCE 80-18770 and 82-15053), the French Centre National de la Recherche Scientifique and the Terres Australes et Antarctiques Françaises, and the National Science Council of the Republic of China. Final writing was supported by NSC (76-0209 M 110-03). We acknowledge assistance of A. POISSON and H. J. MINAS, and valuable comments from two anonymous reviewers.

References

- ACKLEY, S.F., A.J. GOW, K.R. BUCK and K.M. GOLDEN (1980): Sea ice studies in the Weddell Sea aboard USCGC *Polar Sea*. *Antarctic J.*, 84-86.
- ACKLEY, S.F., D.B. CLARKE and S.J. SMITH (1982): Weddell Polynya Expedition Preliminary Data Report: Physical chemical and biological properties of ice cores. Unpublished manuscript.
- ANDERSON, L. and D. DYRSSEN (1980): Constituent data for Leg 2 of the YMER 80 expedition. Univ. of Göteborg, Sweden. 21 pp.
- BRENNECKE, W. (1921): Die ozeanographischen Arbeiten der deutschen antarktischen Expedition 1911-1912. *Arch. deut. Seewarte 1*, Hamburg, 1-216.
- BROECKER, W.S., T. TAKAHASHI and T.-H. PENG (1985): Reconstruction of past atmospheric CO_2 contents from the chemistry of the contemporary ocean: An evaluation. DOE Technical Report, DOE/OR-857. 79 pp.
- BRUJEWIEZ, S.W., A.N. BOGOYAVLENSKY and V.V. MOKIEVSKAYA (1960): Hydrochemical features of the Okhotsk Sea. *Trans. Oceanol. Inst., Acad. Sci. USSR*, **42**, 125-198 (in Russian).
- CHEN, C.T. (1981): Oxygen solubility in seawater. p. 41-55. *In* R. BATTINO (ed.), *Solubility Data Series V. 7, Oxygen and Ozone*. Pergamon Press.
- CHEN, C.T. (1982a): Carbonate chemistry during WEPOLLEX-81. *Antarctic J.*, 1982 Review, 102-103.
- CHEN, C.T. (1982b): On the distribution of anthropogenic CO_2 in the Atlantic and Southern Oceans. *Deep-Sea Res.*, **29**, 563-580.
- CHEN, C.T. (1983): Distributions of dissolved calcium and alkalinity in the Weddell Sea in winter. *Antarctic J.*, 1983 Review, 136-137.
- CHEN, C.T. (1984): Carbonate chemistry of the Weddell Sea. DOE Technical Report, DOE/EV/10611-4. 118 pp.
- CHEN, C.T. (1985): Preliminary observations of oxygen and carbon dioxide of the wintertime Bering Sea marginal ice zone. *Cont. Shelf Res.*, **4**, 465-483.
- CHEN, C.T. and F.J. MILLERO (1979): Gradual increase of oceanic CO_2 . *Nature*, **277**, 205-206.
- CHEN, C.T. and R.M. PYTKOWICZ (1979): On the total CO_2 -titration alkalinity-oxygen system in the Pacific Ocean. *Nature*, **281**, 362-365.
- CHEN, C.T., C.L. WEI and M.R. RODMAN (1985): Carbonate chemistry of the Bering Sea. DOE Technical Report DOE/EV/10611-5. 79 pp.
- CHEN, C.T. and A. POISSON (1986): Indian Ocean gas study. *EOS Oceanogr. Report*, Jan. 4, 1986.
- CHEN, C.T., A. POISSON and C. GOYET (1986): Preliminary data report for the INDIVAT 1 and INDIGO 1/INDIVAT 3 cruises in the Indian Ocean. DOE Technical Report, DOE/NBB-0074. 106 pp.
- CLARKE, D.B. and S.F. ACKLEY (1982): Physical, chemical and biological properties of winter sea ice during the US-USSR Weddell Polynya Expedition. *Antarctic J.*, 1982 Review, 107-109.
- CODISPOTI, L.A., G.E. FRIEDERICK and D.W. HOOD (1986): Variability in the inorganic carbon system over the SE Bering Sea shelf during spring 1980 and spring-summer 1981. *Conti.*

- Shelf Res., **5**, 133-160.
- CSS HUDSON (1984): Cruise 82-001 Data Report, V. 1, Physical and Chemical Data. SIO Reference 84-14. 305 pp.
- DEACON, G.E.R. (1940): Carbon dioxide in the Arctic and Antarctic seas. *Nature*, **145**, 250-252.
- EDMOND, J.M., S.S. JACOBS, A.L. GORDON, A.W. MANTYLA and R.F. WEISS (1979): Water column anomalies in dissolved silica over opaline pelagic sediments and the origin of the deep silica maximum. *J. Geophys. Res.*, **84**, 7809-7826.
- GORDON, A.L., C.T. CHEN and W.G. METCALF (1984): Winter mixed layer entrainment of Weddell Deep Water. *J. Geophys. Res.*, **89**, 637-640.
- GOSINK, T.A., J.G. PEARSON and J.J. KELLEY (1976): Gas movement through sea ice. *Nature* **263**, 41-42.
- HAKODATE MARINE OBSERVATORY (1967): Report of the oceanographic observations in the Tsugaru Straits, sea east of Hokkaido and Tohoku District from February to March 1967. 40 pp.
- HATTORI, A. (1977): Preliminary Report of the Hakuho Maru Cruise KH-75-4. Univ. of Tokyo. 87 pp.
- HATTORI, A. (1979): Preliminary Report of the Hakuho Maru Cruise KH-78-3. Univ. of Tokyo, 87 pp.
- HUBER, B.A., J. JENNINGS, C.T. CHEN, J. MARRA, S. RENNIE, P. MELE and A. GORDON (1983): Reports of the US-USSR Weddell Polynya Expedition, Vol. II. Hydrographical Data LDGO 83-1. 115 pp.
- IGY OCEANOGRAPHY REPORT 2 (1961): Hydrological Observations in the Southern Ocean. 386 pp.
- IVANENKOV, V.I. (1964): Hydrochemistry of the Bering Sea. Akademiya Nauk SSSR, Moscow. 137 pp. (translated by S. Pearson, U.S. Department of Commerce, 1971).
- JENNINGS, J.C., Jr., L.I. GORDON and D.M. NELSON (1984): Nutrient depletion indicates high primary productivity in the Weddell Sea. *Nature*, **308**, 51-54.
- LEVITUS, S. (1982): Climatological atlas of the world ocean. NOAA Professional Paper 13. 173 pp.
- LYAKHIN, Yu. I. and V.P. RUSANOV (1980): Oxygen and carbon dioxide exchange between the Arctic basin and the atmosphere. *Oceanol.*, **20**, 411-418.
- MALMBERG, S.A. (1983): Hydrographic investigations in the Iceland and Greenland Seas in late winter 1971—"Deep Water Project". *Jökull*, **33**, 133-140.
- MINAS, H.J. (1980): Analyse de diagrammes de facteurs hydrologiques et chimiques (temperature, salinite, oxygène, sels nutritifs): Application a l'étude du système production-regeneration dans les resurgences cotières (cotes NW africaines) et les zones à fort melange vertical (Mediterranee, Ocean Antartique). p. 21-36. *In* Production primaire et secondaire, Colloque Fraco-Sovietique, Publ. CNEXO (Actes du colloque) No. 10.
- OB, CRUISE I (1958): Hydrological, hydrochemical, geological and biological studies, Research Ship 'Ob,' 1955-1956. Hydrometeorolog. Publ. House, Leningrad, U.S.S.R.
- OB, CRUISE II (1959): Oceanographic expedition of the Research Ship 'Ob,' 1956-1957. Marine Transport, Leningrad, U.S.S.R.
- POISSON, A. and C.T.A. CHEN (1987): Why is there little anthropogenic CO₂ in the Antarctic Bottom Water? *Deep-Sea Res.*, **34**, 1255-1275.
- REID, J.L. (1973): Northwest Pacific Ocean waters in winter. Johns Hopkins Oceanogr. Studies, No. 5. 96 pp.
- REID, J.L. (1982): On the use of dissolved oxygen concentration as an indicator of winter convection. *NR Reviews*, 28-39.
- SECRETARIA DE MARINA, REPUBLICA ARGENTINA (1959a): Trabajos Oceanograficos Realizados en la Compañia Antartica 1955-1956. Publico H. 620. 137 pp.
- SECRETARIA DE MARINA, REPUBLICA ARGENTINA (1959b): Trabajos Oceanograficos Realizados en la Campaña Antartica 1957-1958. Publico H. 615. 53 pp.
- SECRETARIA DE MARINA, REPUBLICA ARGENTINA (1959c): Trabajos Oceanograficos Realizados en la Campaña Antartica 1958-1959. Publico H. 616. 132 pp.
- SIO REFERENCE 70-5 (1970): Physical, Chemical and Biological Data, Zetes Expedition, Leg 1, 11-24, January 1966. Scripps Inst. Oceanogr. 67 pp.
- SMETANIN, D.A. (1963): Some hydrochemical characteristics of the north-east Pacific based on observations in winter 1958/1959. *Trans. Oceanol. Inst., Acad. Sci. USSR*, **45**, 130-141. (in Russian)
- TAKAHASHI, T., W.S. BROECKER, A.E. BAINBRIDGE and R.F. WEISS (1980): Carbonate chemistry of the Atlantic, Pacific, and Indian Oceans: The results of the GEOSECS expeditions, 1972-1978. Lamont-Doherty Geological Observatory Technical Report No. 1, CV 1-80. 218 pp. (Unpublished document)
- TAKAHASHI, T. and D. CHIPMAN (1982): CO₂ partial pressure in surface waters of the Antarctic Ocean during the US-USSR Weddell

- Polynya Expedition. Antarctic J., 1982 Review 103-104.
- WEISS, R.F., H.G. OSTLUND and H. CRAIG (1979): Geochemical studies of the Weddell Sea. Deep-Sea Res., **26**, 1093-1120.
- WEISS, R.F., W.S. BROECKER, H. CRAIG and D. SPENCER (1983): GEOSECS Indian Ocean Expedition, V. 5, Hydrographic Data 1977-1978. U.S. Government Printing Office, Washington, D.C. 48 pp.

極海中の酸素, 栄養塩, 炭酸塩の夏季と冬季の比較

陳 鎮 東

要旨: 極海における海洋化学的調査は, ごく最近まで主として夏季に行われてきた。作業の困難さもあって, 冬季の化学的データはほとんど得られていなかったが, 世界の海洋の深層底層水は極域で冬季に形成されるから, 極域における冬季のデータの欠如は深層底層水が形成された場所から移動していくにつれてその化学的特性がどのように変化するかを厳密に推定することを困難にしていた。現在では, 冬季のウェッデル海, 南インド洋, ベーリング海において酸素, 栄養塩, アルカリ度, 全炭酸, pH, PCO_2 などのデータが得られている。これらを夏季のデータと比較してみると, 酸素などではその濃度に顕著な差異があるのに対して, アルカリ度などでは同一塩分・同一水温で比較しても顕著な季節変化は認められないことがわかる。